



# What is dementia?

A Guide for Families & Carers

 **etac**<sup>®</sup>

# What is dementia?



Dementia is a syndrome (a group of related symptoms) for a range of progressive conditions that affect the brain. There are over 200 subtypes, most common are: Alzheimer’s disease, vascular dementia, dementia with Lewy bodies, frontotemporal dementia, and mixed dementia.

Dementia damages the nerve cells (neurons) in the brain. Messages can’t be sent to and from the brain effectively, preventing the body from functioning normally.

There are lots of other causes and no two types of dementia are the same. In different types of dementia there is damage to different parts of the brain.

Whatever type of dementia is diagnosed and what part of the brain is affected, each person will experience dementia differently. Dementia can affect a person at any age, but is more commonly diagnosed in people over 65.

Dementia often starts by affecting short-term memory and problems recalling things that happened recently. But dementia can also affect the way people think, speak, perceive things, feel, and behave.

There is a huge amount of social stigma and barriers still associated with dementia and as it progresses, social isolation and loneliness become more likely. Many people with dementia, and their families, become depressed and anxious. This can be because of embarrassment, loss of confidence, lack of appropriate stimulation, routine and purposeful activity.\*

## Other Symptoms experienced

- Difficulties concentrating
- Problems planning, thinking things through
- Struggling with familiar daily tasks, eg following a recipe or using a bank card
- Issues with language and communication, eg trouble remembering the right word or keeping up with a conversation
- Problems judging distances (even though eyesight is fine)
- Aggression, mood changes and difficulties controlling emotions.

## References

This guide has been put together using information from the following sources:

- <https://www.dementiauk.org>
- <https://www.alzheimers.org.uk>
- <https://www.nhs.uk/conditions/dementia/about/>

\* Rachel Hills, Advanced Occupational Therapist, Mental Health Liaison Team

# Common types of dementia



## Alzheimers disease

The most common type of dementia in the UK and accounts for around 60% of diagnoses, although it is comparatively rare for under-65s. It is a physical condition caused by changes in the structure of the brain due to a build-up of proteins. This results in 'plaques' and 'tangles' which damage the brain cells' ability to transmit messages and eventually lead to brain cell death.

Research suggests that changes in the brain can occur up to ten years before a person starts to show symptoms of Alzheimer's disease. The symptoms are usually mild at the beginning and gradually worsen over time.

## Vascular dementia

Vascular dementia is the second most common type of dementia (around 17%). It is caused by problems in the blood supply to the brain cells, commonly due to strokes or a series of small strokes, called Transient Ischaemic Attacks (TIAs), which cause areas of localised cell damage in the brain. Changes in a person's condition because of a stroke or TIA are often sudden, before their condition plateaus.

The signs and symptoms of vascular dementia depend on which area of the brain has been affected. Language, reading, writing and communication can be affected. Memory problems may not be an issue initially, if this area of the brain has not been damaged, but may occur later.

## Symptoms may include

### Alzheimers disease

- Difficulty remembering recent events while having a good memory for past events
- Poor concentration
- Difficulty recognising people or objects
- Poor organisation skills
- Confusion and disorientation
- Slow, muddled, or repetitive speech
- Withdrawal from family and friends
- Problems with decision making, problem solving, planning, and sequencing tasks

### Vascular dementia

- Problems with planning or organising, making decisions or problem solving
- Confusion, disorientation, poor concentration
- Visuospatial skills - problems perceiving objects in three dimensions.
- Problems with reasoning and communication
- Delusions, hallucinations
- Changes in behaviour - agitation, aggressive, depression, disturbed sleep pattern
- Reduced mobility

# Common types of dementia

## Frontotemporal dementia

A progressive condition, so symptoms worsen over time. It tends to affect people aged 45 to 65 years, but they can be older. The areas of the brain most affected are the frontal and temporal lobes.

Frontal lobes are important for personality, social functioning, decision making, problem solving, insight, empathy, motivation, and planning. It can also be called frontal dementia, behavioral variant frontotemporal dementia, or Pick's disease.

Temporal lobes are important for speech, word finding, comprehension and language ability. The hippocampus (important for memory formation) is located in the temporal lobe. This can also be called semantic dementia or primary progressive aphasia

Often frontotemporal dementias are not recognised and diagnosed until the person has had symptoms for four to five years. Early signs are often missed or misattributed by the person, their family and health professionals.

## Lewy body dementia

This form of dementia is a progressive condition caused by protein deposits in the brain. Symptoms worsen over time and it affects movement and motor control. It can also be called dementia with Lewy bodies or Parkinson's disease with dementia (affects about 1/3 of those with Parkinson's Disease).

Memory is often less affected than with other types of dementia, but a person might experience sudden periods of confusion which can change on an hourly basis.

## Mixed dementia

At least one in every ten people with dementia is diagnosed as having more than one type, or mixed dementia. It is much more common in older age groups, such as those over 75. The most common is a combination of Alzheimer's disease and vascular dementia. A mixture of the symptoms associated with the types of dementia they have would be experienced.

## Symptoms may include

### Frontal dementia

- Changes in behaviour and personality
- Apathy
- Obsessive or repetitive behaviours
- Loss of empathy
- Changes in appetite and food eaten
- Difficulties with decision making, problem solving and concentration

### Primary progressive aphasia

- Language difficulties & speech problems
- Reduced comprehension
- Loss of understanding of familiar words difficulty recognising people or objects

### Lewy Body dementia

- Prone to falls
- Tremors (like Parkinson's disease)
- Difficulty swallowing
- Shuffle when walking
- Disrupted sleep patterns due to intense dreams/nightmares
- Visual and auditory hallucinations due to the nerve cell damage.

# Other types of dementia



## Parkinson's Disease

A condition in which a part of the brain called the substantia nigra, loses nerve cells resulting in a reduction of a substance called dopamine, important for regulating body movement. As the disease progresses symptoms can include tremor, shaking, slowness of movement, rigidity, problems with balance, sleep, swallowing, speech and increase the risk of falls. About a third of people with Parkinson's develop dementia, usually in the later stages. This is known as 'Parkinson's disease with dementia' or Lewy body dementia.

## Alcoholic Dementia

Caused by excessive alcohol consumption over a prolonged period. More common in people in their 40s and 50s.

## Huntington's Disease

Huntington's disease is a genetic disorder caused by a faulty gene on chromosome 4. It affects movement but there can be damage to the brain which can develop into dementia.

As the disease progresses it is likely to cause involuntary movements and problems with swallowing, speech, balance and increase the risk of falls. Cognitive processes are also affected such as a decline in thinking and reasoning skills.

## Young onset dementia (YOD)

Defined as dementia diagnosed under the age of 65. It is also referred to as 'early onset' or 'working age' dementia.

### Key Facts\*

- There are currently around 850,000 people with dementia in the UK. This is projected to rise to 1 million by 2025 and 1.6 million by 2040.
- 209,600 people will develop dementia this year – one every 3 minutes
- 1 in 6 people over 80 have dementia
- There are over 42,000 people under 65 with dementia in the UK
- 70% of people in care homes have dementia or severe memory problems.
- Dementia is one of the main causes of disability in later life ahead of cancer, cardiovascular disease, and stroke

### Costs\*

The cost of dementia to the UK is currently £34.7 billion a year; an average annual cost of £32,250 per person with dementia, who with their families pay two-thirds of this cost either in unpaid care or paying for private social care.

Although there is no cure for dementia, scientists and researchers are working hard to find one. Until then, support and treatments are available that can help with symptoms and managing daily life. These can allow people with dementia to lead active, purposeful lives and carry on doing the things that matter to them most.

\* Alzheimer's Society



March 8, 2021

Etac is a world-leading developer and provider of ergonomic assistive devices and patient handling equipment. Our heart lies in the solutions that optimise quality of life for the individual, their family and caregivers.

For the latest news and continuously updated product information – visit: [www.etac.com/uk](http://www.etac.com/uk)

Etac  
Unit D4A, Coombswood Business Park East, Coombswood Way  
Halesowen, West Midlands, B62 8BH  
Tel: 0121 561 2222  
Email: [enquiries@etac.uk.R82.com](mailto:enquiries@etac.uk.R82.com)  
Visit: [www.etac.com/uk](http://www.etac.com/uk)

 **etac**<sup>®</sup>  
Creating Possibilities